

## Operator Algebras Problem set 2

To be submitted by Wednesday, **April 23**, 4 pm.

### Exercise 1 (4 Points)

- Prove that Example 2.3 is indeed a  $C^*$ -algebra, that is, prove that, for any compact Hausdorff space  $X$ ,  $C(X)$  is a commutative unital  $C^*$ -algebra.
- Show that for any two compact Hausdorff spaces  $X$  and  $Y$ , and any continuous mapping  $h : X \rightarrow Y$ , the map  $\alpha_h : C(Y) \rightarrow C(X)$  defined by  $f \mapsto f \circ h$  is a  $*$ -homomorphism.
- Prove that in the situation of (b), if  $h$  is a homeomorphism, the map  $\alpha_h$  is an isometric  $*$ -isomorphism.

### Exercise 2 (4 Points)

Let  $A$  be a non-unital  $C^*$ -algebra. A *double centralizer* of  $A$  is a pair  $(L, R)$  of linear maps  $L, R : A \rightarrow A$  satisfying  $L(ab) = L(a)b$ ,  $R(ab) = aR(b)$  and  $aL(b) = R(a)b$  for all  $a, b \in A$ . The *multiplier algebra*  $M(A)$  of  $A$  is defined as

$$M(A) = \{(L, R) \text{ double centralizer of } A\}$$

with the operations

$$\begin{aligned}(L_1, R_1) + (L_2, R_2) &= (L_1 + L_2, R_1 + R_2) \\ (L_1, R_1) * (L_2, R_2) &= (L_1L_2, R_2R_1) \\ (L, R)^* &= (R^*, L^*) \\ \lambda(L, R) &= (\lambda L, \lambda R)\end{aligned}$$

for  $\lambda \in \mathbb{C}$  and  $(L, R), (L_1, R_1), (L_2, R_2) \in M(A)$ . (Note, for a linear map  $T : A \rightarrow A$ , the map  $T^* : A \rightarrow A$  is defined by  $T^*(x) = (T(x^*))^*$ ). We define a norm on  $M(A)$  via

$$\|(L, R)\| = \|L\| = \|R\|$$

for  $(L, R) \in M(A)$ .

- Show that  $M(A)$  is a unital  $C^*$ -algebra and that the map

$$\phi : A \rightarrow M(A), a \mapsto (L_a, R_a)$$

where  $L_a(x) = ax$  and  $R_a(x) = xa$  for  $x \in A$  is an isometric  $*$ -homomorphism. Furthermore prove that  $\phi(A)$  is an ideal in  $M(A)$ . Thus every non-unital  $C^*$ -algebra can be embedded as an ideal in a unital  $C^*$ -algebra.

- (b) Prove that  $M(A)$  is the largest unitization of  $A$ , that is, if  $B$  is a unital  $C^*$ -algebra and  $A \subset B$  as an ideal, then there exists a  $*$ -homomorphism from  $B$  to  $M(A)$  which extends the embedding  $A \subset M(A)$ .

**Talk** (0 Points)

- (a) Explain that  $A = \bigoplus_{k=1}^N M_{n_k}(\mathbb{C})$  is a  $C^*$ -algebra, where  $n_k \in \mathbb{N}$  and  $M_{n_k}(\mathbb{C})$  is the algebra of  $n_k \times n_k$  matrices.
- (b) Show (or at least motivate) that any finite-dimensional  $C^*$ -algebra is of the form as in (a). This is known as the Theorem of Artin-Wedderburn

Recommended literature: Gerald Murphy “ $C^*$ -algebras and Operator Theory“, Theorem 6.3.8